



Year: 2016

The impact of mild induced hypothermia on the rate of transfusion and the mortality in severely injured patients: a retrospective multi-centre study

Jensen, Kai Oliver ; Held, Leonhard ; Kraus, Andrea ; Hildebrand, Frank ; Mommsen, Philipp ; Mica, Ladislav ; Wanner, Guido A ; Steiger, Peter ; Moos, Rudolf M ; Simmen, Hans-Peter ; Sprengel, Kai

Abstract: BACKGROUND: Although under discussion, induced hypothermia (IH) is an established therapy for patients with cardiac arrest or traumatic brain injuries. The influences on coagulopathy and bleeding tendency in severely injured patients (SIP) with concomitant traumatic brain injury are most widely unclear. Therefore, the aim of this study was to quantify the effect of mild IH in SIP with concomitant severe traumatic brain injuries on transfusion rate and mortality. **METHODS:** In this retrospective multi-centre study, SIP from three European level-1 trauma centres with an ISS 16 between 2009 and 2011 were included. At hospital A, patients qualified for IH with age < 70 years and a severe head injury with an abbreviated injury scale (AISHead) of ≥ 3 . IH was defined as target core body temperature of 35 °C. Hypothermic patients were matched with two patients, one from hospital B and one from hospital C using age and AISHead. The effect of IH on the transfusion rate, complications and mortality was quantified with 95 % confidence intervals (CI). Patients not treated with IH in hospital A and those from hospital B and C, who were not matched, were used to adjust the CI for the effect of inter-hospital therapy protocol differences. **RESULTS:** Mean age of patients in the IH-group (n = 43) was 35.7 years, mean ISS 30 points and sex distribution showed 83.7 % male. Mean age of matched patients in the normotherm-group (n = 86) was 36.7 years, mean ISS 33 points and there were 75.6 % males. For the hypothermic patients, we pointed out an estimate of mean difference for the number of transfused units of packed red blood cells as well as for mortality which does not indicate a decrease in the benefit gained by hypothermia. It is suggested that hypothermic patients tend to a higher rate of lung failure and thromboembolisms. **CONCLUSION:** Though tending to an increased rate of complications, there is no evidence for a difference in both; rate of transfusion and mortality in SIP. Mild IH as an option for severe head injuries seems as well-being practicable in the presence of multiple severe injuries. Further, clinical studies regarding the side effects are necessary.

DOI: <https://doi.org/10.1186/s40001-016-0233-x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-128141>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Jensen, Kai Oliver; Held, Leonhard; Kraus, Andrea; Hildebrand, Frank; Mommsen, Philipp; Mica, Ladislav; Wanner, Guido A; Steiger, Peter; Moos, Rudolf M; Simmen, Hans-Peter; Sprengel, Kai (2016). The impact of mild induced hypothermia on the rate of transfusion and the mortality in severely injured patients: a retrospective multi-centre study. *European Journal of Medical Research*:21:37.
DOI: <https://doi.org/10.1186/s40001-016-0233-x>

RESEARCH

Open Access



The impact of mild induced hypothermia on the rate of transfusion and the mortality in severely injured patients: a retrospective multi-centre study

Kai Oliver Jensen^{1*}, Leonhard Held⁴, Andrea Kraus⁴, Frank Hildebrand², Philipp Mommsen³, Ladislav Mica¹, Guido A. Wanner¹, Peter Steiger⁵, Rudolf M. Moos⁶, Hans-Peter Simmen¹ and Kai Sprengel¹

Abstract

Background: Although under discussion, induced hypothermia (IH) is an established therapy for patients with cardiac arrest or traumatic brain injuries. The influences on coagulopathy and bleeding tendency in severely injured patients (SIP) with concomitant traumatic brain injury are most widely unclear. Therefore, the aim of this study was to quantify the effect of mild IH in SIP with concomitant severe traumatic brain injuries on transfusion rate and mortality.

Methods: In this retrospective multi-centre study, SIP from three European level-1 trauma centres with an ISS ≥ 16 between 2009 and 2011 were included. At hospital A, patients qualified for IH with age ≤ 70 years and a severe head injury with an abbreviated injury scale (AIS_{Head}) of ≥ 3 . IH was defined as target core body temperature of 35 °C. Hypothermic patients were matched with two patients, one from hospital B and one from hospital C using age and AIS_{Head}. The effect of IH on the transfusion rate, complications and mortality was quantified with 95 % confidence intervals (CI). Patients not treated with IH in hospital A and those from hospital B and C, who were not matched, were used to adjust the CI for the effect of inter-hospital therapy protocol differences.

Results: Mean age of patients in the IH-group ($n = 43$) was 35.7 years, mean ISS 30 points and sex distribution showed 83.7 % male. Mean age of matched patients in the normotherm-group ($n = 86$) was 36.7 years, mean ISS 33 points and there were 75.6 % males. For the hypothermic patients, we pointed out an estimate of mean difference for the number of transfused units of packed red blood cells as well as for mortality which does not indicate a decrease in the benefit gained by hypothermia. It is suggested that hypothermic patients tend to a higher rate of lung failure and thromboembolisms.

Conclusion: Though tending to an increased rate of complications, there is no evidence for a difference in both; rate of transfusion and mortality in SIP. Mild IH as an option for severe head injuries seems as well-being practicable in the presence of multiple severe injuries. Further, clinical studies regarding the side effects are necessary.

Keywords: Severely injured, Induced hypothermia, Transfusion rate, Mortality

Background

Hypothermia is classified by causation and extent. Causation of hypothermia can be endogen, controlled-induced

or accidental [1–3]. The extent of hypothermia is mild ($<35-34$ °C), moderate ($34-32$ °C) or severe (<32 °C) [1]. Moderate controlled-induced hypothermia is an established therapy for patients with cardiac arrest in the post-resuscitation phase [4]. Guidelines suggest that a temperature of $32-34$ °C should be applied in pre-hospital as well as in inner-hospital management of cardiac arrest, although Nielsen et al. found out that unconscious

*Correspondence: kaioliver.jensen@usz.ch

¹ Division of Trauma Surgery, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland
Full list of author information is available at the end of the article

survivors after cardiac arrest do not benefit from a targeted temperature of 33 °C in contrast to 36 °C [4–6]. Hypothermia appears to have benefits on elevated intracranial pressure (ICP) in patients with traumatic brain injury within the first 48 h post-trauma if a temperature of 35 °C is applied with external surface cooling or with intra-vascular devices. Patients treated with hypothermia were less likely to die or to have an unfavourable outcome than those in the control group [7, 8]. On the other hand, it is common knowledge that accidentally occurring hypothermia seems to be correlated with a propensity for bleeding complications without being an independent predictor of mortality [9]. Accidental hypothermia, acidosis and coagulopathy represent the lethal triad in severely injured patients [10]. In the current literature, we found no studies in which permissive hypothermia or IH was investigated concerning severely injured patients with concomitant traumatic brain injury. Most studies deal with accidental hypothermia, IH in patients with cardiac arrest or animal trauma models treated with IH [1, 4, 9, 11, 12]. There is thus a lack of information about the effects of IH for severely, multiple injured patients with more than one injured body region. The aim of this study was to highlight the impact of mild IH of 35 °C in these severely injured patients with respect to the transfusion rate within the first 48 h and the in-hospital mortality.

Patients and methods

Study design and hypothesis

We conducted a retrospective multi-centre study to investigate the effect of mild IH in severely injured patients, with a target core temperature of 35 °C on the transfusion rate within the first 48 h post-trauma and on in-hospital mortality in comparison to matched normotherm patients.

Patients

All patient data were prospectively collected for the internal trauma databases from three European University level-1 trauma centres, hereafter called hospital A, hospital B and hospital C. Inclusion criteria were severely injured patients, defined by an ISS ≥ 16 including severe head injury, defined by an AIS_{Head} ≥ 3 , admission within the period from January 2009 to December 2011 and need for an initial intensive care treatment after damage control surgery [13]. Excluded were patients with restricted comfort therapy due to declared intention, age <18 years, an early relocation within the first 48 h and a late admission of ≥ 4 h counted from time of injury.

Methods

Cooling

In hospital A, hypothermia is induced using a clinically approved algorithm for traumatic brain injuries, although

other injuries may be present (Fig. 1). The intended temperature for hypothermia in hospital A was 35 °C. Down-regulation of the temperature was accomplished by a regime with different phases: (1) removal of blanket; (2) Coolmat (Blanketrol® II Hyper-Hypothermia System, Cincinnati Sub-Zero Products Inc., Cincinnati, USA); (3) Coolgard® (Femoral or subclavian vein) (Zoll Cool-Line catheter, Zoll Circulation Inc., Sunnyvale, USA). The minimum duration of hypothermia was specified as 24 h. Within the protocol of hypothermia, deep analgesation was performed. In hospitals B and hospital C, the aim of temperature management was normothermia. Within this study none of the patients had to be cooled via Coolgard®.

Measurement of body temperature


The temperature was measured in a standardised manner depending on availability. If available a probe for continuous monitoring of brain tissue oxygen and brain temperature was installed (Neurovent-P-Temp, Raumedic AG, Helmbrechts, Germany). Following a descending order, the ocular surface temperature, measurements via pulse contour cardiac output (PiCCO) (Picco 2, Pulsion Medical Systems SE, Feldkirchen, Germany) or aural temperature probe were used.

Definition of organic failure

Organic failure (OF) is defined as at least one organ scoring three or more points according to the Sequential Organ Failure Assessment (SOFA) score, matching the following criteria: *cardio circulatory system* dopamine greater than 5 µg/kg per minute or any adrenaline/noradrenaline treatment; *lung* Horowitz score (PaO₂/FiO₂) of less than 200 mmHg; *liver* serum bilirubin level of ≥ 6.0 mg/dL; *kidney* serum creatinine levels of 3.5 mg/dL or greater or urine output of less than 500 cc/day; *CNS* Glasgow Coma Scale (GCS) of 8 points or less [14]; multi-organ failure (MOF) is defined by the simultaneous failure of two or more organ systems. Sepsis is defined by the ACCP/SCCM Consensus Conference criteria [9, 15].

Variables

Primary outcome variables were the number of transfused units of packed red blood cells (EC) and mortality. In addition, we recorded number of transfused units of fresh-frozen plasma (FFP) and thrombocyte units (TC) in cc. Secondary outcome variables such as organic failure (OF) distinguished by relevant organs, MOF, the occurrence of sepsis and thromboembolism (TE) were recorded as well. For the primary outcome variables, a length of stay (LOS) in hospital of 48 h or more was required, whilst for secondary outcome variables, an



LOS in hospital of at least 7 days without relocation was necessary.

Matching and statistical analysis

During the evaluation period, in hospital A hypothermia was induced as a standard of care in patients when aged ≤ 70 years and suffering from a severe head injury with an abbreviated injury scale (AIS_{Head}) of ≥ 3 . In consecutive absence of an inner-hospital control group, we matched patients treated with IH with one patient of hospital B and one patient from hospital C according to the characteristics, that determined the qualification for hypothermia in hospital A [16].


We estimated the effect of IH by means of a 95 % confidence interval (CI). For continuous outcomes (e.g. EC, FFP, TC), we computed non-parametric bootstrap CIs for the mean difference between the outcomes of the matched patients [17]. For binary outcomes (e.g. death), we constructed Wald CI for the log odds ratio, using conditional logistic regression. To explore the difference in outcome due to inter-hospital therapy protocol differences and to reduce the risk of a possible selection bias, we estimated the hospital effect with 95 % CI using the outcomes of patients who were not treated with IH in hospital A as well as for patients from hospitals B and hospital C, who were not matched to the patients treated with IH in hospital A. In this analysis, we adjusted for the age and ISS of the patient. Finally, the estimated hospital effect was subtracted from the estimated effect of hypothermia computed in the first step, and the two CIs

were combined using the square-and-add approach, to obtain CI for the effect of IH [18]. Confidence intervals for log odds ratios were transformed to the risk difference scale assuming a baseline mortality equal to the average mortality of the patients from hospitals B and hospital C matched to the patients from hospital A. The analysis was performed separately for the two hospital pairs (A vs. B and A vs. C). Statistical analysis has been performed in the R programming language [19].

Results

A total of 1808 patients were evaluated in this study. Regarding the underlying inclusion and exclusion criteria 564 patients were included, of whom 43 patients were treated with hypothermia in hospital A. A data set was created with 129 patients including the 43 hypothermic patients in hospital A as well as the matched normothermic patients from hospital B and hospital C ($n = 86$) (Fig. 2).

The mean ISS of the hypothermic patients was 30 points (SD 9.6 points), mean age was 35.7 years (SD 13.2 years) with 83.7 % males. The mean ISS of the matched normothermic patients was 33 points (SD 12.4 points), mean age was 36.7 years (SD 13.3 years) and 75.6 % patients were males. Base data for each hospital are shown in Table 1 separately. AIS distribution regarding the main six regions of interest (head; face; thorax; abdomen; extremities; external) is demonstrated in Table 2. In mean, 60.5 % of the hypothermic and matched patients suffered from a relevant injury (AIS ≥ 3) of other regions than AIS_{Head}.



Primary outcomes

Considering the rate of EC transfused within the first 48 h post-trauma, an estimated mean of -1.5 units [95 % CI -5.3 to 2.5 for hospital A in comparison to hospital B (AB)] and -1.3 units (95 % CI -3.3 to 0.8 AC) was calculated. We found an RD for mortality of -0.14 (95 % CI -0.29 to 0.14 AB) and -0.18 (95 % CI -0.30 to 0.07 for AC) (Figs. 3 and 4).

Secondary outcomes

The existence of IH seems to have a negative side effect on the prevalence of organic failure of the lungs with OR of 4.7 (95 % CI 1.0 – 22.6 AB) and 4.7 (95 % CI 0.9 – 24.4 AC) and the occurrence of thromboembolism with OR of 12.9 (95 % CI 1.2 – 133.9 AB) and of 21.0 (95 % CI 2.0 – 217.4 AC) (Table 3).

Due to the small number of specific complications, the OR for sepsis as well as for organic failure of

liver and kidneys was not further taken into account. Organic failure of the heart and the central nervous system are secondary outcomes, which cannot be used for further comparison due to the hypothermia protocol used in hospital A, including the proposed deep analgesedation with a consecutive Glasgow Coma Scale of 3 and the common application of noradrenaline leading to an organic failure classification (central nervous system, heart) following the SOFA score [14]. The variable multi-organ failure defined as a combination of two or more organic failures is falsified as a sequella.

Discussion

The present study indicates that mild IH of 35°C neither has a negative effect on the coagulation system of severely injured patients measured by transfusion rate nor on the mortality. Actual literature suggests a benefit of 15 % for

Table 1 Base data separately for each hospital divided into hypotherm, matched normotherm and remaining cases of normotherm population

	Hospital					
	A	B	C	A	B	C
	Hypotherm	Matched	Matched	Normotherm		
Variable	<i>n</i> = 43	<i>n</i> = 43	<i>n</i> = 43	<i>n</i> = 275	<i>n</i> = 97	<i>n</i> = 63
Age (years)	36 ± 13	38 ± 13	36 ± 14	55 ± 22	50 ± 23	61 ± 18
ISS	30 ± 10	33 ± 10	33 ± 14	28 ± 10	27 ± 8	27 ± 10
NISS	44 ± 14	43 ± 13	43 ± 18	37 ± 13	32 ± 12	36 ± 14
TRISS	0.68 ± 0.28	0.73 ± 0.26	0.64 ± 0.36	0.71 ± 0.31	0.69 ± 0.31	0.61 ± 0.33
Blunt trauma mechanism (%)	97.6	95.1	100	95.6	97.9	98.4
Sex female (%)	16.3	27.9	18.6	33.8	34.0	25.4
At arrival in resuscitation room						
SAP (mmHg)	130 ± 26	115 ± 32	116 ± 40	135 ± 32	124 ± 31	127 ± 42
Heart rate (beats/min)	87 ± 22	94 ± 25	82 ± 31	87 ± 22	86 ± 23	90 ± 25
Temperature (°C)	35.3 ± 1.2	35.1 ± 1.3	34.7 ± 2.4	35.2 ± 1.4	35.4 ± 1.2	35.6 ± 2.1
Haemoglobin (g/L)	12.5 ± 1.8	11.6 ± 2.7	13.1 ± 2.2	12.1 ± 2.5	11.5 ± 2.6	12.0 ± 2.9
Thrombocytes (10 ³ /μL)	218 ± 57	194 ± 60	216 ± 72	224 ± 80	198 ± 62	214 ± 73
Prothrombin time (%)	80 ± 18	71 ± 25	71 ± 25	74 ± 25	72 ± 21	78 ± 20
Base excess (mmol/L)	−3.3 ± 2.8	−3.4 ± 5.2	−2.6 ± 4.2	−2.4 ± 4.7	−2.6 ± 3.7	−3.3 ± 4.3
Lactate (mmol/L)	2.3 ± 1.7	2.8 ± 2.0	3.6 ± 3.7	2.2 ± 1.8	2.6 ± 1.8	2.9 ± 2.6

Data given as mean and standard deviation, ISS injury severity score, NISS new ISS, TRISS trauma ISS, SAP systolic arterial pressure

induced hypothermia [20]. Our data showed an upper margin of the 95 % CI within this range for mortality and the comparison between both hospital pairs. The same evidence was given for the amount of EC transfused. We could not prove an increase in bleeding complications or in-hospital mortality due to hypothermia.

Underlining our conclusions, an improvement of survival was measured in an uncontrolled haemorrhagic shock model comparing moderate hypothermia versus normothermia [21, 22]. Whilst another study ascribes significant coagulopathy in severely injured patients to hypothermia, it has only been shown for hypothermia below 34 °C. The authors presume hypercoagulability to partially compensate the restrictions of coagulation due to hypothermia [23]. These results do conform to the findings in the present study with an aim of 35 °C. In the same way, the mortality due to accidental hypothermia varies from 30 % up to 100 % depending on the depth of accidental hypothermia [2, 24, 25]. Accidental hypothermia seems to trigger the vicious circle of the lethal triad of hypothermia, acidosis and coagulopathy [10].

Based on our results, we agree with Mohr et al. who reported of an IH animal trauma model without an increase of coagulopathy. Within that study, there were no findings showing negative impact of hypothermia on inflammatory complications and mortality [12]. Other authors mention the negative effects of hypothermia without further exploring them [26].



Although our investigations suggested lung failure and TE to be bigger problems for patients treated with hypothermia, it is not associated with a higher mortality but only with morbidity. Related to these findings, Alderson et al. showed that hypothermia resulted in a statistically significant increase in the OR for incidence of pneumonia and a decrease in the OR for death in patients with head injuries, whereas Geurts et al. observe in a meta-analysis no increase in the overall risk of pneumonia or sepsis but strongly suggest an association between them [27, 28]. Descending into the matter of blood levels of TNF-alpha and IL-6 in hypothermic animals as inflammatory cytokine mediators, Gröger et al. found almost similar values at 35 °C and with normothermia in

Table 2 AIS distribution by hospital for hypothermic and normothermic matched patients

AIS region	Value	Hospital		
		A (n)	B (n)	C (n)
Head	3	3	3	3
	4	20	19	14
	5	20	21	26
Face	0	22	28	23
	1	4	6	5
	2	9	6	8
	3	8	3	6
	4	0	0	1
	5	0	0	0
Thorax	6	0	0	0
	0	21	9	25
	1	0	2	1
	2	4	10	6
	3	14	17	5
	4	4	3	4
	5	0	2	1
Abdomen	6	0	0	1
	0	31	29	33
	1	0	0	0
	2	11	12	1
	3	1	1	5
	4	0	1	4
	5	0	0	0
Extremities	6	0	0	0
	0	25	18	27
	1	0	1	1
	2	11	11	8
	3	6	9	5
	4	1	4	1
	5	0	0	1
External	6	0	0	0
	0	22	35	33
	1	14	6	4
	2	6	2	5
	3	1	0	0
	4	0	0	1
	5	0	0	0
	6	0	0	0

contrast to hypothermia of 32 °C [29]. In summary, the temperature seems to be mainly affecting the morbidity of lung failure and TE. Even though we expected an association between IH and morbidity, we could not show an increase of in-hospital mortality.

We are aware of some weaknesses in this study. The main handicap is the retrospective nature of the study,



although all data were recorded prospectively. Additionally, a larger number of matched cases would provide greater confidence in the significance of the results. Furthermore, the absence of an inner-hospital control group due to the cooling-protocol in hospital A lead to this elaborate statistical design.

We conclude that severely injured patients did not suffer from relevant bleeding complications due to IH because of a similar transfusion rate.

Table 3 95 % confidence intervals for all outcomes

Outcome parameter	Comparison	Estimate	95 % CI from	To
EC difference	AB	−1.5	−5.3	2.5
	AC	−1.3	−3.3	0.8
FFP difference	AB	1.2	−2.0	4.0
	AC	0.7	−1.7	2.8
TC difference	AB	−0.5	−0.9	0.3
	AC	−0.1	−0.3	0.2
Death (OR)	AB	0.5	0.1	1.8
	AC	0.4	0.1	1.3
Of lung (OR)	AB	4.7	1.0	22.6
	AC	4.7	0.9	24.4
TE (OR)	AB	12.9	1.2	133.9
	AC	21.0	2.0	217.4

EC number of transfused units of packed red blood cells, FFP number of transfused units of fresh-frozen plasma, TC number of transfused units of thrombocyte units, OF organ failure, TE thromboembolism, CI confidence interval, AB hospital A in comparison to hospital B, AC hospital A in comparison to hospital C, OR odds ratio

Conclusion

The combination of the potential positive effects of hypothermia for traumatic brain injuries and of not increasing the usage of blood products or the mortality makes mild controlled-induced hypothermia a feasible procedure for severely, multiple injured patients with concomitant, severe traumatic brain injury. We look forward to upcoming prospectively randomised controlled studies on this topic with great pleasure and interest.

Abbreviations

IH: induced hypothermia; SIP: severely injured patients; ISS: injury severity score; AIS: abbreviated injury scale; CI: confidence interval; PiCCO: pulse contour cardiac output; SOFA: Sequential Organ Failure Assessment; GCS: Glasgow Coma Scale; MOF: multi-organ failure; EC: units of packed red blood cells; FFP: units of fresh-frozen plasma; OF: organ failure; TE: thromboembolism; LOS: length of stay; TC: thrombocyte units; SD: standard deviation; RD: risk difference or RISC difference; OR: odds ratio; cCT: cranial computed tomography; paCO₂: arterial carbon dioxide pressure; CPP: cerebral perfusion pressure; S_{jv}O₂: jugular venous oxygen saturation; ptiO₂: brain tissue oxygen; TCD: transcranial doppler.

Authors' contributions

KS conceived and together with KOJ and FH designed the study. FH, PS and PM were involved in data collection. LH and AK performed the statistical analysis and preparation of figures. KOJ drafted the paper. LM, GAW, RMM and HPS contributed substantially to its revision. All authors read and approved the final manuscript.

Author details

¹ Division of Trauma Surgery, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland. ² Department of Orthopedic Trauma, University of Aachen, Aachen, Germany. ³ Trauma Department, Hannover Medical School, Hannover, Germany. ⁴ Department of Biostatistics, Institute for Epidemiology, Biostatistics and Prevention, University of Zurich, Zurich, Switzerland. ⁵ Division of Surgical Intensive Care Medicine, University Hospital Zurich, University of Zurich, Zurich, Switzerland. ⁶ University Hospital Zurich, University of Zurich, Zurich, Switzerland.

Acknowledgements

We would like to thank H. Ingleby for editing and proofreading the manuscript and S. Angst for basic data management.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The datasets during and/or analysed datasets during the current study are available from the corresponding author on reasonable request.

Compliance with ethical requirements

Approval by the local ethics committee was obtained (KEK-ZH-Nr. 2013-0023) and this study has therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. Due to this ethical approval no consent to participate in the study was needed, cause only anonymised data were used.

Received: 29 July 2016 Accepted: 30 September 2016

Published online: 06 October 2016

References

- Hildebrand F, Probst C, Frink M, Huber-Wagner S, Krettek C. Importance of hypothermia in multiple trauma patients. *Der Unfallchirurg*. 2009;112(11):959–64. doi:10.1007/s00113-009-1683-1.
- Hildebrand F, Giannoudis PV, van Griensven M, Chawda M, Pape HC. Pathophysiologic changes and effects of hypothermia on outcome in elective surgery and trauma patients. *Am J Surg*. 2004;187(3):363–71. doi:10.1016/j.amjsurg.2003.12.016.
- Seekamp A, Hildebrand F, van Griensven M, Grotz M, Tscherne H. Role of adenosine triphosphate (ATP) in trauma-induced and elective hypothermia. *Zentralbl Chir*. 1999;124(11):1017–29.
- Kim F, Nichol G, Maynard C, Hallstrom A, Kudenchuk PJ, Rea T, et al. Effect of prehospital induction of mild hypothermia on survival and neurological status among adults with cardiac arrest: a randomized clinical trial. *JAMA*. 2014;311(1):45–52. doi:10.1001/jama.2013.282173.
- Werdan K, Russ M, Buerke M, Delle-Karh G, Geppert A, Schöndube FA. Cardiogenic shock due to myocardial infarction: diagnosis, monitoring and treatment: a German-Austrian S3 Guideline. *Dtsch Arzteblatt Int*. 2012;109(19):343–51. doi:10.3238/arztebl.2012.0343.
- Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, et al. Targeted temperature management at 33 degrees C versus 36 degrees C after cardiac arrest. *N Engl J Med*. 2013;369(23):2197–206. doi:10.1056/NEJMoa1310519.
- Sydenham E, Roberts I, Alderson P. Hypothermia for traumatic head injury. *Cochrane Database Syst Rev*. 2009; doi:10.1002/14651858.CD001048.pub3.
- Sydenham E, Roberts I, Alderson P. Hypothermia for traumatic head injury. *Cochrane Database*. 2009. doi:10.1002/14651858.CD001048.pub4.
- Trentzsch H, Huber-Wagner S, Hildebrand F, Kanz KG, Faist E, Piltz S, et al. Hypothermia for prediction of death in severely injured blunt trauma patients. *Shock (Augusta, Ga)*. 2012;37(2):131–9. doi:10.1097/SHK.0b013e318245f6b2.
- Mikhail J. The trauma triad of death: hypothermia, acidosis, and coagulopathy. *AACN Clin Issues*. 1999;10(1):85–94.
- Seekamp A, Ziegler M, Van Griensven M, Grotz M, Regel G. The role of hypothermia in trauma patients. *Euro J Emer Med*. 1995;2(1):28–32.
- Mohr J, Ruchholtz S, Hildebrand F, Flohe S, Frink M, Witte I, et al. Induced hypothermia does not impair coagulation system in a swine multiple trauma model. *J Trauma Acute Care Surg*. 2013;74(4):1014–20. doi:10.1097/TA.0b013e3182826edd.
- Taeger G, Ruchholtz S, Waydhas C, Lewan U, Schmidt B, Nast-Kolb D. Damage control orthopedics in patients with multiple injuries is effective, time saving, and safe. *J Trauma*. 2005;59(2):409–16.
- Vincent JL, Moreno R, Takala J, Willatts S, De Mendonca A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the working group on

- sepsis-related problems of the European society of intensive care medicine. *Intensive Care Med.* 1996;22(7):707–10.
15. American College of Chest Physicians/Society of Critical Care. Medicine Consensus Conference: definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med.* 1992;20(6):864–74.
 16. Rosenbaum PR. *Design of Observational Studies*. New York: Springer; 2012.
 17. Davison AC, Hinkley DV. *Bootstrap methods and their application* Cambridge series in statistical and probabilistic mathematics. Cambridge: Cambridge University Press; 2003.
 18. Newcombe RG. *Confidence intervals for proportions and related measures of effect size*. London: Taylor & Francis; 2012.
 19. R Development Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2009.
 20. Andrews PJ, Sinclair LH, Harris B, Baldwin MJ, Battison CG, Rhodes JK, et al. Study of therapeutic hypothermia (32 to 35 degrees C) for intracranial pressure reduction after traumatic brain injury (the Eurotherm3235Trial): outcome of the pilot phase of the trial. *Trials.* 2013;14:277. doi:[10.1186/1745-6215-14-277](https://doi.org/10.1186/1745-6215-14-277).
 21. Kentner R, Rollwagen FM, Prueckner S, Behringer W, Wu X, Stezoski J, Safar P, Tisherman SA. Effects of mild hypothermia on survival and serum cytokines in uncontrolled hemorrhagic shock in rats. *Shock.* 2002;17(6):521–6.
 22. Takasu A, Carrillo P, Stezoski SW, Safar P, Tisherman SA. Mild or moderate hypothermia but not increased oxygen breathing prolongs survival during lethal uncontrolled hemorrhagic shock in rats, with monitoring of visceral dysoxia. *Crit Care Med.* 1999;27(8):1557–64.
 23. Watts DD, Trask A, Soeken K, Perdue P, Dols S, Kaufmann C. Hypothermic coagulopathy in trauma: effect of varying levels of hypothermia on enzyme speed, platelet function, and fibrinolytic activity. *J Trauma.* 1998;44(5):846–54.
 24. Tsuei BJ, Kearney PA. Hypothermia in the trauma patient. *Injury.* 2004;35(1):7–15.
 25. Jurkovich GJ, Greiser WB, Luteran A, Curreri PW. Hypothermia in trauma victims: an ominous predictor of survival. *J Trauma.* 1987;27(9):1019–24.
 26. Kobbe P, Lichte P, Wellmann M, Hildebrand F, Nast-Kolb D, Waydhas C, et al. Impact of hypothermia on the severely injured patient. *Der Unfallchirurg.* 2009;112(12):1055–61. doi:[10.1007/s00113-009-1711-1](https://doi.org/10.1007/s00113-009-1711-1).
 27. Alderson P, Gadkary C, Signorini DF. Therapeutic hypothermia for head injury. *Cochrane Database Syst Rev.* 2004. doi:[10.1002/14651858.CD001048.pub2](https://doi.org/10.1002/14651858.CD001048.pub2).
 28. Geurts M, Macleod MR, Kollmar R, Kremer PH, van der Worp HB. Therapeutic hypothermia and the risk of infection: a systematic review and meta-analysis. *Critical Care Med.* 2014;42(2):231–42. doi:[10.1097/CCM.0b013e3182a276e8](https://doi.org/10.1097/CCM.0b013e3182a276e8).
 29. Groger M, Scheuerle A, Wagner F, Simon F, Matallo J, McCook O, et al. Effects of pretreatment hypothermia during resuscitated porcine hemorrhagic shock. *Critical Care Med.* 2013;41(7):e105–17. doi:[10.1097/CCM.0b013e31827c0b1f](https://doi.org/10.1097/CCM.0b013e31827c0b1f).

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

